

Two Families of Quantum Codes Derived from Cyclic Codes

K. Guenda

Faculty of Mathematics, University of Sciences and Technology of Algiers

B. P 32 El Alia 16111 Bab Ezzouar, Algiers, Algeria

guendakenza@hotmail.com

Abstract

We characterize the affine-invariant maximal extended cyclic codes. Then by the *CSS* construction, we derive from these codes a family of pure quantum codes. Also for $\text{ord}_n q$ even, a new family of degenerate quantum stabilizer $[[n, 1, \geq \sqrt{n}]]_q$ codes is derived from the classical duadic codes. This answer an open problem asked by Aly et al.

MSC:[94B05, 94B15, 94B60]

Keywords: Cyclic codes, affine-invariant codes, duadic codes, CSS construction, quantum codes.

1 Introduction

In this paper we consider maximal-affine invariant codes and duadic codes. The first are extended cyclic codes and the second are cyclic codes. We characterize the form of the maximal-affine invariant codes and the cases when they contain their dual codes. Then by using the CSS construction we derive a new family of quantum codes. For q a prime number, the obtained quantum codes are pure. In the second part of this paper, the interesting relation between the duals of the pairwise of even-like and odd-like duadic codes allow us to give a $[[n, 1, \geq d]]_q$ family of quantum degenerate codes. The degeneracy of these codes is an interesting propriety as it was shown in [1]. The construction given here is more general than in [1], since it does not require that $\text{ord}_n q$ is odd. Note that the quantum codes are used to protect quantum information over noisy quantum channels.

2 The Maximal Affine-Invariant Codes

Let $q = p^r$ be a power of a prime number p , $K = \mathbb{F}_{p^r}$ and $G = \mathbb{F}_{p^m}$ be respectively the finite field with p^r and p^m elements and $m = rm'$. A primitive cyclic code C of length $n = p^m - 1$ is an ideal of the algebra $R = \mathbb{F}_q[x]/(x^n - 1)$ generated by a polynomial $g(x)$ which divides $x^n - 1$, and is uniquely determined by its defining set $T = \{0 \leq i \leq n-1 \mid g(\beta^i) = 0\}$, where β is an n^{th} primitive root of the unity in G . The set T is then an union of cyclotomic classes $Cl(j) = \{jq^l \bmod n \mid l \in \mathbb{Z}\}$. If T contains one cyclotomic class, C

is called maximal cyclic code. We associate with the code C the code

$$\hat{C} = \{(x_0, \dots, x_n) \mid (x_1, \dots, x_n) \in C \text{ with } x_0 = -\sum_{i=1}^n x_i\},$$

called the extended code of C . The defining set T of C is also called the defining set of $\hat{C}$. Since the code $\hat{C}$ is of length p^m , the field G is viewed as the support field and the coordinate position of the codewords are labelled by the elements of G , hence can be considered as a subspace of the group algebra $A = K[\{G, +\}]$, which is the group algebra of the additive group of G over K ,

$$A = \left\{ \sum_{g \in G} x_g X^g, x_g \in K \right\}.$$

The operations in A are given by

$$a \sum_{g \in G} x_g X^g + b \sum_{g \in G} y_g X^g = \sum_{g \in G} (ax_g + by_g) X^g, \quad a, b \in K,$$

and

$$\sum_{g \in G} x_g X^g \times \sum_{g \in G} y_g X^g = \sum_{g \in G} \left(\sum_{h+k=g} x_h y_k \right) X^g.$$

The zero and unity of A are $\sum_{g \in G} 0 X^g$ and X^0 .

The permutation group $\text{Per}(C)$ of a linear code of length n is the subgroup of $\text{Sym}(n)$ which leaves the code C invariant. An extended cyclic code $\hat{C}$ of length p^m is said to be affine-invariant provided its permutation group contains $\text{AGL}(1, p^m) = \{\sigma \mid f_\sigma(X) = aX + b, a \in G^*, b \in G\}$.

A partial order on $S = [0, p^m - 1[$ is defined by:

If $s = \sum_{i=0}^{m-1} s_i p^i$, $t = \sum_{i=0}^{m-1} t_i p^i$ are the p -adic expansions of s and t , then

$$s \prec t \iff s_i \leq t_i, \forall i = 0, \dots, m-1, s_i \in [0, p-1].$$

The following well known result [3, 6] characterizes the affine-invariant codes by their defining sets.

Theorem 1 An extended cyclic code $\hat{C}$ with defining set T is affine-invariant if and only if T satisfies

$$\forall t \in T, \forall s \in S, s \prec t \Rightarrow s \in T.$$

As a consequence we get the following result.

Proposition 2 A maximal extended cyclic code of A is affine-invariant if and only if

$$T = Cl(p^j).$$

Proof. Let T be the defining set of a maximal extended cyclic code, it is clear that if $T = Cl(p^j)$ with $0 \leq j \leq m-1$, then for $s \in S$ and $t \in T$ we can have $s \prec t$ only for $s = t$. Therefore by Theorem 1 the set T is the defining set of an affine-invariant code.

Let $T = Cl(t)$ be a cyclotomic class modulo $p^m - 1$, where t is the smallest

element for the order ($\leq$). For $t \in]p^j, p^{j+1}[$, the p -adic expansion of t is

$$t = \sum_{i=0}^{j-1} s_i p^i + \alpha p^j,$$

with $0 \leq s_i < p$ and $1 \leq \alpha$. Since $1 \leq \alpha$, then $p^j \prec t$, and by Theorem 1 the extended code associated to T cannot be affine-invariant. ■

Remark 3 Using the fact that $\text{ord}_{p^m-1} p^r = \frac{m}{r}$, we can obviously deduce that the cardinality of $Cl(p^j)$ is $\frac{m}{r}$. Hence for fixed parameters, the number of distinct maximal affine-invariant codes is r .

3 Quantum Affine-Invariant Codes

From the classical affine-invariant codes of the previous section we can directly obtain a family of quantum codes by using the called CSS given by the following Lemma.

Lemma 4 Let $C_1 = [n, k_1, d_1]$ and $C_2 = [n, k_2, d_2]$ be linear codes over $\mathbb{F}_q$ with $C_2^\perp \subset C_1$. Furthermore, let

$$d_Q = \min\{wt(v) : v \in (C_1 \setminus C_2^\perp) \cup (C_2 \setminus C_1^\perp)\} \geq \min(d_1, d_2).$$

Then there exists a $[[n, k_1 + k_2 - n, d_Q]]_q$ quantum code.

Proof. See for instance [4]. ■

By Lemma 4, if a linear $[n, k, d]$ code C contains its dual $C^\perp$, then there exists a quantum code with parameters

$$[[n, 2k - n, d_Q \geq d]]_q,$$

that is pure up to d . Now we prove the following Lemma which tell us when a maximal affine-invariant code can contains its dual.

Lemma 5 Let $\hat{C}$ be an extended maximal affine-invariant code $[p^m, p^m - 1 - \frac{m}{r}, d]$, then if $p > 3$ or $m > 2$ or $r \neq 1$ we have $\hat{C}^\perp \subset \hat{C}$.

Proof. Let $T^\perp$ be the defining set of $\hat{C}^\perp$, we have the relation $T^\perp = \{s \in \mathbb{F}_{p^m} \mid n - s \notin T\}$. To have $\hat{C}^\perp \subset \hat{C}$ it suffices to have $T \subset T^\perp$, which is then equivalent to the following statement

$$s \in T \Rightarrow p^m - 1 - s \notin T. \quad (1)$$

Let $T = Cl(p^j)$, for simplicity we consider $s = p^j$; the general case is therefore obviously deduced. Seeking a contradiction we assume that $p^m - 1 - p^j \in T$, then it exists $0 \leq \alpha \leq \frac{m}{r} - 1$ such that $p^m - 1 - p^j = p^{\alpha r + j} \pmod{p^m - 1}$, this is equivalent to have $p^j(1 + p^{\alpha r}) = k(p^m - 1)$ for some $k \geq 1$. Hence

$1 + p^{\alpha r} = k'(p^m - 1)$ for some $k' \geq 1$, this implies that

$$1 + p^{\alpha r} \geq p^m - 1. \quad (2)$$

Since $\alpha \leq \frac{m}{r} - 1$, then (2) gives $2 \geq p^{m-r}(p^r - 1)$, this is possible only for $(r = m = 1, p \leq 3)$ or $(r = 1, m = 2, p = 2)$. Therefore under the hypothesis $p > 3$ or $m > 2$ or $r \neq 1$ we have $\hat{C}^\perp \subset \hat{C}$. ■

From Lemma 5 and the CSS construction we get the following result.

Theorem 6 Let $q = p^r$ with p a prime number, m a positive integer and $n = p^m - 1$. If $p > 3$ or $m > 2$ or $r \neq 1$, then there exists a quantum code with parameters

$$[[p^m, p^m - 2 - 2\frac{m}{r}, \frac{m}{r} + 2 \geq d_Q \geq d_A]]_q,$$

where d_A is the minimum distance of an extended maximal affine-invariant code.

Proof. Proposition 2 and Lemma 5 imply the existence of affine-invariant code $\hat{C}$ which contains its dual with parameters $[p^m, p^m - 1 - \frac{m}{r}, d_A]$. Therefore by Lemma 4, we have the existence of a $[[p^m, p^m - 2 - 2\frac{m}{r}, d_Q \geq d_A]]_q$ quantum code. The quantum Singleton bound gives : $d_Q \leq 2 + \frac{m}{r}$. ■

Corollary 7 Assuming that $r = 1$ and $(p > 3$ or $m > 2)$, there exists a

pure quantum code with parameters

$$[[p^m, p^m - 2 - 2m, m + 2 \geq d_Q = d_A]]_p,$$

where d_A is the minimum distance of an extended maximal affine-invariant code.

Proof. Assume $r = 1$ and $(p > 3$ or $m > 2)$, then Theorem 6 gives the existence of a quantum code with parameters $[[p^m, p^m - 2 - 2m, m + 2 \geq d_Q \geq d_A]]_p$. From Remark 3 there is a unique maximal affine invariant code $\hat{C}$, which is the extended BCH code with designed distance 2. Then from the Weil bound given in [12] the minimum distance of $\hat{C}^\perp$ is $d = p^m - p^{m-1}$ or $d = p^m - p^{m-1} + 1$. Hence from Lemma 4 the quantum code is pure; $d_Q = d_A$.

■

4 Duadic Codes

Throughout this part we denote by n an odd integer and by q a prime power. The notation $q \equiv \square \pmod n$ express the fact that q is a quadratic residue modulo n . We write $p^\alpha \parallel m$ if and only if the integer m is divisible by p^α but not by $p^{\alpha+1}$. For an integer a such that $(a, n) = 1$, $\mu_a : i \longmapsto ia \pmod n$ denote a permutation on $Z_n = \{0, \dots, n-1\}$.

4.1 Classical Duadic Codes

Let q be a prime power and n an odd integer such that $(n, q) = 1$. Let S_1 and S_2 be unions of cyclotomic cosets modulo n such that $S_1 \cap S_2 = \emptyset$, $S_1 \cup S_2 = Z_n \setminus \{0\}$ and $aS_i \bmod n = S_{(i+1) \bmod 2}$. Then the triple μ_a, S_1, S_2 is called splitting modulo n . The odd-like duadic codes D_1 and D_2 are the cyclic codes over $\mathbb{F}_q$ with defining set respectively S_1 and S_2 . The even-like duadic codes C_1 and C_2 are the cyclic codes over $\mathbb{F}_q$ with defining set respectively $\{0\} \cup S_1$ and $\{0\} \cup S_2$.

Theorem 8 Duadic codes of length n over $\mathbb{F}_q$ exist if and only if

$$q \equiv \square \pmod{n}.$$

Proof. This is well-known, see for example [11]. ■

Lemma 9 (square root bound)

Let D_1 and D_2 be a pair of odd-like duadic codes of length n over $\mathbb{F}_q$. Then they have the same minimum distance say d , and verifies

1. $d^2 \geq n$
2. $d^2 - d + 1 \geq n$ if the splitting is given by μ_{-1}

Proof. See [8, Theorem 7]. ■

4.2 Quantum Duadic Codes

If we are over $\mathbb{F}_{q^2}$, we can consider the Hermitian duality. Furthermore we have the existence of duadic codes over $\mathbb{F}_{q^2}$ for all n , when $(n, q^2) = 1$, since $q^2 \equiv \square \pmod{n}$.

The following Lemma gives the same construction as Lemma 4 in the Hermitian case.

Lemma 10 *If there exists a classical linear $[n, k, d]_{q^2}$ code C , such that $C \subset C^{\perp h}$. Then there exists an $[[n, n - 2k, \geq d]]_q$ quantum code that is pure to d .*

Proof. See for instance [4, Corollary 2]. ■

Lemma 11 *Let C_i and D_i respectively be the even-like and odd-like codes over $\mathbb{F}_{q^2}$, where $i \in \{0, 1\}$. Then $C_i^{\perp h} = D_i$ if and only if there is a q^2 -splitting given by μ_{-q} , that is, $-qS_i \equiv S_{(i+1 \bmod 2)} \pmod{n}$.*

Proof. See [9, Theorem 4.4] ■

We prove the following Lemma which gives some properties of $\text{ord}_n q$ which are useful for our construction.

Lemma 12 *Let q be a prime power and n an integer, then :*

1. *If $\text{ord}_n q$ is odd, then $\text{ord}_n q^2 = \text{ord}_n q$.*
2. *If $\text{ord}_n q$ is even, then $\text{ord}_n q^2 = \frac{\text{ord}_n q}{2}$*

Proof.

1. Let $r = \text{ord}_n q$ and $r' = \text{ord}_n q^2$, then $r|2r'$, hence $r|r'$ since r is odd. On the other hand, we have $q^{2r} \equiv 1 \pmod n$, therefore $r'|r$, finally $r = r'$.
2. Let $r = \text{ord}_n q$ and $r' = \text{ord}_n q^2$, we have $q^{2r'} \equiv 1 \pmod n$ which implies that $r|2r'$. Since r is even, then $q^{2\frac{r}{2}} \equiv 1 \pmod n$ therefore $r'|\frac{r}{2}$, hence $r' = \frac{r}{2}$.

■

Proposition 13 Let $n = \prod p_i^{m_i}$ be the prime factorization of the odd integer n , where each $m_i > 0$, such that every $p_i \equiv -1 \pmod 4$, then $\text{ord}_n q^2$ is odd.

Proof. We have $\text{ord}_n q^2 = \text{lcm}(\text{ord}_{p_i^{m_i}} q^2)$. By [7] we have that $\text{ord}_{p_i^{m_i}} q^2 = p_i^{m_i - z_i} t_i$, where $t_i = \text{ord}_{p_i} q^2$ and z_i is such that $p_i^{z_i} \parallel q^{t_i} - 1$. Hence to prove the proposition it suffices to prove that $\text{ord}_{p_i} q^2$ is odd if $p_i \equiv -1 \pmod 4$. We have to consider two cases:

1. $q \equiv \square \pmod{p_i}$, then $\text{ord}_{p_i} q$ divides $\frac{p_i-1}{2}$, this implies that $\text{ord}_{p_i} q$ is odd otherwise $p_i \equiv 1 \pmod 4$. Then from Lemma 12 $\text{ord}_{p_i} q^2$ is odd.
2. q is not a square modulo p_i , then $\text{ord}_{p_i} q = p_i - 1$ is even and then from Lemma 12 $\text{ord}_{p_i} q^2 = \frac{p_i-1}{2}$. The last quantity must be odd otherwise we get $p_i \equiv 1 \pmod 4$.

■

Lemma 14 Let $n = \prod p_i^{m_i}$ be an odd integer such that $\text{ord}_n q^2$ is odd. Then μ_{-q} gives a splitting of n over $\mathbb{F}_{q^2}$. Furthermore μ_{-1} and μ_{-q} give the same splitting.

Proof. Let $\{S_1, S_2, a\}$ be a splitting over $\mathbb{F}_{q^2}$. We have $q^{\text{ord}_n q^2} S_i = S_i \pmod n$. If $\text{ord}_n q^2 = 2k + 1$, then $q^{2k+1} S_i = q^{2k} q S_i = S_i$, but since we are over $\mathbb{F}_{q^2}$ we have $q^{2k} S_i = S_i$, hence $q S_i = S_i$. That means that μ_q fixes each S_i if the multiplicative order of q^2 modulo n is odd. From [9, Lemma 5], there exists a q^2 -splitting of n given by μ_{-1} if and only if $\text{ord}_n q^2$ is odd. Hence $-S_i \equiv S_{(i+1 \bmod 2)} \pmod n$; since μ_q fixes S_i hence μ_{-q} gives a q^2 -splitting of n . Conversely if μ_{-q} gives a q^2 -splitting of n , then $-q S_i \equiv S_{(i+1 \bmod 2)} \pmod n$. Since μ_q fixes S_i we have $-q S_i \equiv -S_i \equiv S_{(i+1 \bmod 2)} \pmod n$; hence μ_{-q} gives the same splitting as μ_{-1} . ■

Now by using the results of Lemma 14, we construct a quantum family codes. This construction is more general than the ones done in [1, Theorem 10].

Theorem 15 Let n be an odd integer such that $\text{ord}_n q^2$ is odd. Then there exists an $[[n, 1, d]]_q$ quantum code with $d^2 - d + 1 \geq n$.

Proof. From Lemma 14 there exist duadic codes $C_i \subset D_i$ with splitting given by μ_q and μ_{-1} . This means that the $C_i \subset C_i^{\perp h} = D_i$, by Lemma 11. Therefore by Lemma 10 there exists an $[[n, n - (n - 1), d]]_q$ quantum code with $d = \text{wt}(D_i \setminus C_i)$. Since μ_{-1} gives a splitting, from Lemma 9 we have $d^2 - d + 1 \geq n$. ■

Now by using Theorem 15 we construct a family of degenerate codes.

Theorem 16 Let $n = \prod p_i^{m_i}$ be an odd integer such that every $p_i \equiv -1 \pmod 4$. Let $t_i = \text{ord}_{p_i} q^2$ and $p_i^{z_i} \parallel q^{2t_i} - 1$. Then for $m_i > 2z_i$, there

exists degenerate $[[n, 1, d]]_q$ quantum codes pure to $d' \leq \min\{p_i^{z_i}\} < d$ with $d^2 - d + 1 \geq n$.

Proof. From [1, Lemma 5] there exists an even-like duadic codes with parameters $[n, (n-1)/2, d']_{q^2}$ and $d' \leq \min\{p_i^{z_i}\}$. Then by [11, Theorem 8] the splitting for these codes is given by μ_{-1} . By Proposition 13 $\text{ord}_n q^2$ is odd and then by Lemma 14 μ_{-q} also gives a splitting for these codes. Hence by Theorem 16 this duadic code gives a quantum duadic code $[[n, 1, d]]_q$, which is impure as $d' \leq \min\{p_i^{z_i}\} < \sqrt{n} < d$. ■

References

- [1] S. A. Aly, A. Klappenecker and P. K. Sarvepalli, Remarkable degenerate quantum stabilizer codes derived from duadic codes, *arXiv: quant-ph/0601117v1*, Jan 2006.
- [2] T. P. Berger and P. Charpin, The automorphism groups of *BCH* codes and of some affine-invariant codes over extension fields, *Design, Codes and Cryptography*, 18, 29-53, 1999.
- [3] P. Charpin, Codes Cycliques étendus invariants sous le groupe affine, *Thèse de Doctorat d'État Université de Paris VII*, Jan 1987.
- [4] M. Grassl, T. Beth and M. Rötteler, On optimal quantum codes, *Internat. J. Quantum Information*, 2(1):757-775, 2004.
- [5] W. C. Huffman and V. Pless, Fundamentals of error-correcting codes. *Cambridge*, 2003.

- [6] T. Kasami, S. Lin and W. W. Peterson, Some results on cyclic codes which are invariant under the affine group and their application, *Inform and Control*, 11: 475-496, 1967.
- [7] W. J. LeVeque, Elementary number theory of numbers. *Dover, New York*, 1990.
- [8] V. Pless Duadic Codes Revisited, *Congressus Numerantium*, 59: 225-233, 1987.
- [9] J. J. Rushanun, Topics in Integral Matrices and Abelian Group Codes, *Ph.D. Thesis, California Institute of Technology*, 1986.
- [10] J. J. Rushanun, Duadic codes and difference sets, *J. Combin. Theory Ser. A* 57: 254-261, 1991.
- [11] M. H. M. Smid, Duadic codes, *IEEE Trans. Inform*, 29(2), 1983.
- [12] J. Wolfman, New bounds on cyclic codes from algebraic curves , in “*Lecture Notes in Computer Science*”, 388: 47-62, *Springer-Verlag* 1989.